


# Trigonometric Ratios



Suppose  $\theta$  is any angle in standard position, and P (x,y) is any point on its terminal arm, at a distance of r from the origin. The value of r can then be determined using the Pythagorean Theorem,  $r = \sqrt{x^2 + y^2}$

## PRIMARY TRIGONOMETRIC RATIOS

The three primary trigonometric ratios can be defined in terms of x, y, and r as follows:

$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{y}{r} \quad \cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{x}{r} \quad \tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{y}{x}$$

## RECIPROCAL TRIGONOMETRIC RATIOS

The three reciprocal trigonometric ratios can be defined in terms of x, y, and r as follows:


$$\text{cosecant } \theta = \frac{1}{\sin \theta} \quad \text{secant } \theta = \frac{1}{\cos \theta} \quad \text{cotangent } \theta = \frac{1}{\tan \theta}$$

OR, abbreviated and in terms of x, y, and r:

$$\csc \theta = \frac{r}{y} \quad \sec \theta = \frac{r}{x} \quad \cot \theta = \frac{x}{y}$$

## THE CAST RULE

The six trigonometric ratios of any angle in the *first* quadrant are always positive, however, this is not the case in the other quadrants. For each quadrant, we will determine the sign for each of the *primary* trigonometric ratios and summarize the results with the CAST rule.



We can determine the six trigonometric ratios for any angle in standard position using:

- i. the coordinates of the point where terminal arm intersects the unit circle, and/or
- ii. the special triangles

Example 1: Determine the Trigonometric Ratios for Angles in the Unit Circle

The point  $A\left(\frac{-12}{13}, \frac{5}{13}\right)$  lies at the intersection of the unit circle and the terminal arm of an angle  $\theta$  in standard position.

- a. Draw a diagram to model the situation.
- b. Determine the values of the six trigonometric ratios for  $\theta$ . Express your answers in lowest terms.

Solution:

<p>a.</p> 	<p>b.</p>
--	-----------

Example 2: Exact Values for Trigonometric Ratios

*Exact values for the trigonometric ratios can be determined by using the unit circle or special triangles.*

Determine the exact value for each trigonometric ratio.

a. $\sin \frac{5\pi}{6}$	b. $\cos \frac{-2\pi}{3}$
c. $\csc 315^\circ$	d. $\tan 180^\circ$

### Example 3: Approximate Values of Trigonometric Ratios

You can determine approximate values for sine, cosine and tangent using a calculator. Remember to set your calculator to either degree or radian setting, depending on the question. To find the value of a trigonometric ratio for cosecant, secant or cotangent, use the appropriate reciprocal relationship.

For example,  $\csc 4.1 = \frac{1}{\sin 4.1} = -1.2220$  (Set calculator to radians)

Determine the approximate value for each trigonometric ratio. Give your answers to four decimal places.

a. $\tan \frac{9\pi}{5}$	b. $\sin 220^\circ$	c. $\cos 1.25$	d. $\sec(-110^\circ)$

### Example 4: Evaluating Trigonometric Ratios

Determine the *exact* value for each of the following trigonometric expressions.

a.  $\sin 45^\circ \cos 45^\circ + \sin 30^\circ \sin 60^\circ$

b. 
$$\frac{2\sin^2 \frac{3\pi}{4} + \cos^2 \frac{5\pi}{6}}{\cos \frac{2\pi}{3}}$$

c. 
$$\frac{3\cos 180^\circ + \sin 135^\circ}{\sin 30^\circ}$$

**Example 5: Calculating Trigonometric Values for Points Not on the Unit Circle**

The point  $A(6, -8)$  lies on the terminal arm of an angle  $\theta$  in standard position.

- What is the exact value of each trigonometric ratio for  $\theta$ ?
- Determine  $\theta$  in the domain  $-4\pi \leq \theta \leq 4\pi$ .

**Solution:**

**Example 6: Find Angles Given Their Trigonometric Ratios**

Determine the measures of all angles that satisfy the following.

- $\cos \theta = 0.598472$  in the domain  $0 \leq \theta < 2\pi$ . Give your answers to the nearest tenth of a radian.
- $\sin \theta = -0.819152$  in the domain  $0^\circ \leq \theta < 360^\circ$ . Give your answers to the nearest degree.
- $\cos \theta = \frac{-\sqrt{2}}{2}$  in the domain  $0 \leq \theta < 4\pi$ . Give exact answers.
- $\tan \theta = \frac{1}{\sqrt{3}}$  in the domain  $-180^\circ \leq \theta < 180^\circ$ . Give exact answers.
- $\csc \theta = -\frac{2}{\sqrt{3}}$  in the domain  $-2\pi \leq \theta < \pi$ . Give exact answers.

**Solution:**

- $\cos \theta = 0.598472$  in the domain  $0 \leq \theta < 2\pi$ . Give your answers to the nearest tenth of a radian.

b.  $\sin \theta = -0.819152$  in the domain  $0^\circ \leq \theta < 360^\circ$ . Give your answers to the nearest degree.

c.  $\cos \theta = \frac{-\sqrt{2}}{2}$  in the domain  $0 \leq \theta < 4\pi$ . Give exact answers.

d.  $\tan \theta = \frac{1}{\sqrt{3}}$  in the domain  $-180^\circ \leq \theta < 180^\circ$ . Give exact answers.

e.  $\csc \theta = -\frac{2}{\sqrt{3}}$  in the domain  $-2\pi \leq \theta < \pi$ . Give exact answers.